

From AMS Glossary

radar equation

An equation expressing the power of a radar echo at the input of the receiving antenna of a radar as a function of the range and radar cross section of a target.

For a point target and plane-polarized radiation, it is written

$$P_r = P_t \frac{G^2 \lambda^2 \sigma}{64 \pi^3 r^4},$$

where P_r is the received power, P_t the peak power transmitted by the radar, G the gain of the antenna, λ the wavelength, r the distance to the target, and σ the radar cross section of the target. For a distributed target such as precipitation, which fills the radar beam, the radar equation may be written

$$\overline{P_r} = P_t \frac{G^2 \lambda^2 \theta \phi h}{1024 \pi^2 \ln 2} \frac{\eta}{r^2},$$

where θ and ϕ denote the antenna beamwidths in the horizontal and vertical planes, h is the pulse length of the transmitted signal, and η is the radar reflectivity of the target. This equation assumes that the antenna pattern has a Gaussian shape and that the scattering volume is uniformly filled. The radar signals received from distributed targets fluctuate; the overbar on P_r indicates that it is a time average over a period equal to several multiples of the coherence time of the received signal. From the radar equation, the fundamental measurable property of precipitation is the radar reflectivity η , which depends on the sizes and concentration of the hydrometeors and their thermodynamic phase. For hydrometeors small enough for the Rayleigh scattering approximation, it is given by

$$\eta = \frac{|K|^2 \pi^5}{\lambda^4} Z,$$

where

$|K|^2$ is a dielectric factor, approximately equal to 0.93 for water and 0.21 for ice. The factor Z is called the radar reflectivity factor of the precipitation. It equals the sum of the sixth-powers of the diameters of the water drops in a unit volume of space or of the melted diameters of the snow and ice particles in a unit volume. It may be expressed in terms of the drop-size distribution as

$$Z = \int_0^{\infty} N(D) D^6 dD,$$

where $N(D) dD$ is the number of drops per unit volume with diameters in the interval dD . (For ice-phase precipitation, $N(D)$ is the distribution of melted diameters.)

Retrieved from "http://glossary.ametsoc.org/w/index.php?title=Radar_equation&oldid=17790"

Categories: Term | Terms R

-
- This page was last modified on 25 April 2012, at 19:42.

- This page has been accessed 5,446 times.